

002065



CENTRAL LIBRARY  
MAHIDOL UNIVERSITY

PRODUCTION OF CHOLESTEROL OXIDASE FOR  
CLINICAL ASSAY OF CHOLESTEROL

BY

VIMONRUTH SUPITUG

(B.Sc. in Medical Technology)

A THESIS SUBMITTED IN PARTIAL FULFILLMENT OF  
THE REQUIREMENT FOR THE DEGREE OF  
MASTER OF SCIENCE

IN

BIOCHEMISTRY

FACULTY OF GRADUATE STUDIES

OF

MAHIDOL UNIVERSITY

1987

Copyright by Mahidol University

อภินันท์นากการ

จาก

บัณฑิตวิทยาลัย

มหาวิทยาลัย

ABSTRACT

Cholesterol determination is routinely performed in hospitals as a diagnostic test for some pathological state such as arteriosclerosis or jaundice etc. In the present time determination of cholesterol by chemical reagents has been replaced by enzymatic reagents. The assay system requires cholesterol esterase, cholesterol oxidase, and peroxidase. As a result the enzymatic method which is superior to chemical methods in all aspect is rather expensive. The major objective of this project was to find suitable source of cholesterol oxidase so that enzymic reagent for cholesterol determination could be produced locally.

Screening for the enzyme source was carried out in various kind of vegetable and in microorganism. Cholesterol oxidase was found in one of 276 isolates of actinomycetes. The morphology and culturing condition of the isolate has been investigated.

Evaluation of the suitability of the enzyme produced from this isolate of actinomycete was carried out by partial purification of the enzyme, using DEAE-cellulose chromatography and gel filtration. The partially purified enzyme was studied for its physical and biochemical properties, as followings : molecular weight is 34,000,  $K_m$  is  $1.47 \times 10^{-4} M$ , the enzyme is specific for  $3\beta$ -hydroxysterols with double bond at C-5, stability to heating is up to  $45^\circ C$ , stability at  $-10^\circ C$  for one month with 5% loss of activity, optimum pH is around 6.5 and optimum temperature is at  $40^\circ C$ .

The partially purified enzyme was compared with commercial cholesterol oxidase in the determination of serum cholesterol. Correlation ( $r = .987$ ) between the two source of enzyme was statistically significant.

From this study it was concluded that cholesterol oxidase by actinomycete is very suitable in all aspects to be used as a diagnostic enzyme.

